



Nakajo-Nishimura syndrome

Nakajo-Nishimura syndrome is an inherited condition that affects many parts of the body and has been described only in the Japanese population. Beginning in infancy or early childhood, affected individuals develop red, swollen lumps (nodular erythema) on the skin that occur most often in cold weather; recurrent fevers; and elongated fingers and toes with widened and rounded tips (clubbing).

Later in childhood, affected individuals develop joint pain and joint deformities called contractures that limit movement, particularly in the hands, wrists, and elbows. They also experience weakness and wasting of muscles, along with a loss of fatty tissue (lipodystrophy), mainly in the upper body. The combination of muscle and fat loss worsens over time, leading to an extremely thin (emaciated) appearance in the face, chest, and arms.

Other signs and symptoms of Nakajo-Nishimura syndrome can include an enlarged liver and spleen (hepatosplenomegaly), a shortage of red blood cells (anemia), a reduced amount of blood clotting cells called platelets (thrombocytopenia), and abnormal deposits of calcium (calcification) in an area of the brain called the basal ganglia. Intellectual disability has been reported in some affected individuals.

The signs and symptoms of Nakajo-Nishimura syndrome overlap with those of two other conditions: one called joint contractures, muscular atrophy, microcytic anemia, and panniculitis-induced lipodystrophy (JMP) syndrome; and the other called chronic atypical neutrophilic dermatosis with lipodystrophy and elevated temperature (CANDLE) syndrome. All three conditions are characterized by skin abnormalities and lipodystrophy. Although they are often considered separate disorders, they are caused by mutations in the same gene, and some researchers believe they may represent different forms of a single condition.

Frequency

Nakajo-Nishimura syndrome appears to be rare and has been described only in the Japanese population. About 30 cases have been reported in the medical literature.

Genetic Changes

Nakajo-Nishimura syndrome is caused by mutations in the *PSMB8* gene. This gene provides instructions for making one part (subunit) of specialized cell structures called immunoproteasomes, which are found primarily in immune system cells. Immunoproteasomes play an important role in regulating the immune system's response to foreign invaders, such as viruses and bacteria. One of the primary functions of immunoproteasomes is to help the immune system distinguish the body's

own proteins from proteins made by foreign invaders, so the immune system can respond appropriately to infection.

Mutations in the *PSMB8* gene greatly reduce the amount of protein produced from the *PSMB8* gene, which impairs the normal assembly of immunoproteasomes and causes the immune system to malfunction. For unknown reasons, the malfunctioning immune system triggers abnormal inflammation that can damage the body's own tissues and organs; as a result, Nakajo-Nishimura syndrome is classified as an autoinflammatory disorder.

Abnormal inflammation likely underlies many of the signs and symptoms of Nakajo-Nishimura syndrome, including the nodular erythema, recurrent fevers, joint problems, and hepatosplenomegaly. It is less clear how mutations in the *PSMB8* gene lead to muscle wasting and lipodystrophy. Studies suggest that the protein produced from the *PSMB8* gene may play a separate role in the maturation of fat cells (adipocytes), and a shortage of this protein may interfere with the normal development and function of these cells.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- ALDD
- autoinflammation, lipodystrophy, and dermatosis syndrome
- Japanese autoinflammatory syndrome with lipodystrophy
- JASL
- Nakajo syndrome
- NKJO

Diagnosis & Management

These resources address the diagnosis or management of Nakajo-Nishimura syndrome:

- Genetic Testing Registry: Nakajo syndrome
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1850568/>

These resources from MedlinePlus offer information about the diagnosis and management of various health conditions:

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Clubbed Fingers (image)
<https://medlineplus.gov/ency/imagepages/18127.htm>
- Encyclopedia: Immune Response
<https://medlineplus.gov/ency/article/000821.htm>
- Health Topic: Fever
<https://medlineplus.gov/fever.html>
- Health Topic: Skin Conditions
<https://medlineplus.gov/skinconditions.html>

Genetic and Rare Diseases Information Center

- Nakajo Nishimura syndrome
<https://rarediseases.info.nih.gov/diseases/3916/nakajo-nishimura-syndrome>

Additional NIH Resources

- National Institute of Arthritis and Musculoskeletal and Skin Diseases:
Autoinflammatory Diseases
https://www.niams.nih.gov/Health_Info/Autoinflammatory/

Educational Resources

- Cambridge University Hospital (UK): Lipodystrophy
http://www.cuh.org.uk/sites/default/files/publications/PIN3418_lipodystrophy_v1.pdf
- Disease InfoSearch: Nakajo syndrome
<http://www.diseaseinfosearch.org/Nakajo+syndrome/5100>

- MalaCards: autoinflammation, lipodystrophy, and dermatosis syndrome
http://www.malacards.org/card/autoinflammation_lipodystrophy_and_dermatosis_syndrome
- Orphanet: Nakajo-Nishimura syndrome
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=2615

Patient Support and Advocacy Resources

- Autoinflammatory Alliance
<http://autoinflammatory.org/candle.php>
- Resource List from the University of Kansas Medical Center: Dermatology and Genetics
<http://www.kumc.edu/gec/support/derm.html>
- Stop Childhood Autoinflammatory Diseases Now
<http://www.stopcaidnow.org/>

Genetic Testing Registry

- Nakajo syndrome
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1850568/>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22autoinflammation%2C+lipodystrophy%2C+and+dermatosis+syndrome%22>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28nakajo+syndrome%5BTIAB%5D%29+OR+%28nakajo-nishimura+syndrome%5BTIAB%5D%29%29+OR+%28%28PSMB8%5BTIAB%5D%29+AND+%28lipodystrophy%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

OMIM

- AUTOINFLAMMATION, LIPODYSTROPHY, AND DERMATOSIS SYNDROME
<http://omim.org/entry/256040>

Sources for This Summary

- Arima K, Kinoshita A, Mishima H, Kanazawa N, Kaneko T, Mizushima T, Ichinose K, Nakamura H, Tsujino A, Kawakami A, Matsunaka M, Kasagi S, Kawano S, Kumagai S, Ohmura K, Mimori T, Hirano M, Ueno S, Tanaka K, Tanaka M, Toyoshima I, Sugino H, Yamakawa A, Tanaka K, Niikawa N, Furukawa F, Murata S, Eguchi K, Ida H, Yoshiura K. Proteasome assembly defect due to a proteasome subunit beta type 8 (PSMB8) mutation causes the autoinflammatory disorder, Nakajo-Nishimura syndrome. *Proc Natl Acad Sci U S A*. 2011 Sep 6;108(36):14914-9. doi: 10.1073/pnas.1106015108. Epub 2011 Aug 18.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21852578>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3169106/>
- Kanazawa N. Nakajo-Nishimura syndrome: an autoinflammatory disorder showing pernio-like rashes and progressive partial lipodystrophy. *Allergol Int*. 2012 Jun;61(2):197-206. doi: 10.2332/allergolint.11-RAI-0416. Epub 2012 Mar 25. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22441638>
- Kitamura A, Maekawa Y, Uehara H, Izumi K, Kawachi I, Nishizawa M, Toyoshima Y, Takahashi H, Standley DM, Tanaka K, Hamazaki J, Murata S, Obara K, Toyoshima I, Yasutomo K. A mutation in the immunoproteasome subunit PSMB8 causes autoinflammation and lipodystrophy in humans. *J Clin Invest*. 2011 Oct;121(10):4150-60. doi: 10.1172/JCI58414. Epub 2011 Sep 1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21881205>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3195477/>
- Kitano Y, Matsunaga E, Morimoto T, Okada N, Sano S. A syndrome with nodular erythema, elongated and thickened fingers, and emaciation. *Arch Dermatol*. 1985 Aug;121(8):1053-6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/4026345>
- Kunimoto K, Kimura A, Uede K, Okuda M, Aoyagi N, Furukawa F, Kanazawa N. A new infant case of Nakajo-Nishimura syndrome with a genetic mutation in the immunoproteasome subunit: an overlapping entity with JMP and CANDLER syndrome related to PSMB8 mutations. *Dermatology*. 2013;227(1):26-30. doi: 10.1159/000351323. Epub 2013 Aug 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23942189>
- Tanaka M, Miyatani N, Yamada S, Miyashita K, Toyoshima I, Sakuma K, Tanaka K, Yuasa T, Miyatake T, Tsubaki T. Hereditary lipo-muscular atrophy with joint contracture, skin eruptions and hyper-gamma-globulinemia: a new syndrome. *Intern Med*. 1993 Jan;32(1):42-5. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/8495043>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/nakajo-nishimura-syndrome>

Reviewed: November 2013

Published: February 14, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services